



Presentation:

Basic

Image:

Small

Français



(1 of 1)

[View Images](#)

PUBLISHED INTERNATIONAL APPLICATION

- (11) **WO 98/03675** (13) A1
(21) **PCT/US97/13321**
(22) **23 July 1997 (23.07.1997)**
(25) ENG (26) ENG
(31) 60/022,354 (32) **24 July 1996** (33) US
(24.07.1996)
(43) 29 January 1998 (29.01.1998)
(51)⁶ C12Q 1/60
(54) CHOLESTEROL SEPARATION AND FLUORESCENT ANALYSIS
(71) **HELENA LABORATORIES CORPORATION** 1530 Lindbergh Drive, P.O. Box 752, Beaumont, TX 77704-0752 ; (US). [US/US]. (*for all designated states except US*)
(72)(75) **HICKS, Debra, Linn** 4233 Holland Avenue, Orange, TX 77630 ; (US) [US/US].
MERCHANT, Mark, Edwin 3316 Omaha, Nederland, TX 77627 ; (US) [US/US].
GUADAGNO, Philip, Angelo 465 Concord, Vidor, TX 77662 ; (US) [US/US].
ROBINSON, Suzan, Sha 126 Inwood, Silsbee, TX 77656 ; (US) [US/US]. **MILLICAN, Stacey, Eloise** 1033 Sunmeadow, Beaumont, TX 77706 ; (US) [US/US]. **NAKAZATO, Tokiya** 9-21-19, Tokiwa, Urawa-shi, Saitama 336 ; (JP) [JP/JP].
(74) **SCHNEIDER, Jerold, I.** Dorsey & Whitney LLP, Suite 200, 1330 Connecticut Avenue N.W., Washington, DC 20036 ; (US).
(81) JP, US ; EP (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE)

No Image Available.

Abstract

A method and reagent for cholesterol fraction separation by electrophoresis and quantitative interpretation of the HDL, LDL and VLDL fractions. The reagent is applied after the electrophoretic separation and each fraction will fluoresce in response to excitation at a wavelength which peaks at 356nm. The reagent includes NAD (nicotinamide adenine dinucleotide) which, in the reduced form NADH, will fluoresce.



Presentation:

Basic

Image:

Small

Français



(1 of 1)